

GLUTEUS MEDIUS/MINIMUS PARTIAL THICKNESS REPAIR POST OPERATIVE PROTOCOL

	Time Frame (Weeks)	Guidelines	Goals
PHASE I	0 to 4	<p>WEIGHT BEARING: 20% FFWB x 4 weeks BRACE: Hip brace set 30-75 of hip flexion x4 weeks ROM: Flexion, extension and IR as tolerated; No ER or Adduction x 4 wks</p> <p>EXERCISE PROGRESSION</p> <ul style="list-style-type: none"> • Lie on stomach 2 or more hours a day • Stationary bike with no resistance immediately as tolerated • Glute, quadriceps, hamstring isometrics (2x/day): Immediately as tolerated (may initiate BFR as available) • Hip PROM (2x/day) flexion, abduction, IR supine at 90° and prone at 0° • Hip circumduction • Initiate basic core: pelvic tilting, TVA/breathing re-education • Quadriceps Stretching in prone • SAQ/LAQ as sitting tolerance improves • Prone assisted hip extension with pillow under hips beginning POD 14 • Quadruped rocking beginning POD 28 <p>MANUAL INTERVENTION</p> <ul style="list-style-type: none"> • Scar mobilization • STM to quad, ITB, hip flexors, glutes, hip adductors/abductors/rotators • Continue work on ROM as outlined above 	<ul style="list-style-type: none"> • Reduce inflammation • Decrease pain <p>POOL PROGRAM:</p> <ul style="list-style-type: none"> • May begin deep water pool walking (chest deep) at 1 week if incisions well covered with tegaderm <p>PHASE II PROGRESSION CRITERIA:</p> <ul style="list-style-type: none"> • Mobility within limitations • Early restoration of neuromuscular control • 6 week Follow-up Exam with surgeon
PHASE II	4 to 10	<p>EXERCISE PROGRESSION</p> <p>Beginning week 4</p> <ul style="list-style-type: none"> • Double leg bridge • Prone hip extension • Mini squat • Standing hip abduction (surgical leg only) • Double leg heel raise • Prone IR/ER AAROM → AROM • Front plank on knees <p>Beginning week 6</p> <ul style="list-style-type: none"> • Bridge progressions (modified single leg bridge foot on chair, bridge march) • Bilateral standing hip abduction (no resistance) • Quadruped hip extension bilateral • Chair squat • Small step up • Clamshell at 60 deg hip flexion (no resistance) • Reverse clamshell (no resistance) • Front plank on toes (no LE lifts) <p>Beginning week 8</p> <ul style="list-style-type: none"> • Standing hip abduction (band above knees) • Clamshell at 30 deg hip flexion (no resistance) • Single leg bridge variations <p>MANUAL INTERVENTION</p> <ul style="list-style-type: none"> • Scar mobilization • STM to quad, ITB, hip flexors, glutes, hip adductors/abductors/rotators • Continue work on ROM (FABER, flexion, abduction, IR, ER) • LE stretching program (avoiding ITB and Piriformis) 	<ul style="list-style-type: none"> • Begin PWB as tolerated with goal to wean off crutches (1-2 wk process) • Normal gait • Normal single limb stance • Full ROM • Improve LE muscle activation, strength and endurance <p>POOL PROGRAM:</p> <ul style="list-style-type: none"> • May begin swimming with pool buoy at 8 weeks <p>PHASE III PROGRESSION CRITERIA</p> <ul style="list-style-type: none"> • Flexion, ER and IR ROM within normal limits • Normal Gait • No Trendelenberg with Single Leg Stance/descending stairs • Normal bilateral squat

	Time Frame (Weeks)	Guidelines	Goals
PHASE III	10 to 20	<p>EXERCISE PROGRESSION</p> <ul style="list-style-type: none"> • Standing hip abduction band moving distally • Side plank progressions knees —> toes • Clamshell/reverse clamshell variations with resistance • Sidelying hip abduction • Quadruped and standing fire hydrant variations • Continue with muscle activation series (quadruped or straight leg series) • Introduce movement series to increase proprioception, balance, and functional flexibility • Progress core program as appropriate including unilateral front planks, side plank clamshell and hip abduction variations • Advanced glute and posterior chain strengthening including hip hikes • Leg press and leg curl • Squat progression (double to single leg: load as tolerated) • Lunge progression • Step-up Progression • Walking program • May swim using flutter/dolphin kick at 10 weeks • Outdoor biking- discuss with surgeon and PT • Implement full LE stretching program as tolerated <p>CV EXERCISE:</p> <ul style="list-style-type: none"> • May begin elliptical and stair climber at 12 weeks • May begin return to run program if phase 4 criteria are met (~6 months) <p>MANUAL INTERVENTION</p> <ul style="list-style-type: none"> • Continue soft tissue mobilization as needed particularly glutes, adductors, hip flexors, abduc-tors • Gentle joint mobilizations as needed for patients lacking ER or FABER ROM • May begin trigger point dry needling for glutes, quads, adductors NO HIP FLEXOR TDN until Week 8. • Assess FMA and begin to address movement dysfunctions 	<p>PHASE IV PROGRESSION CRITERIA</p> <ul style="list-style-type: none"> • 5-6 months post-op • Hip abduction and extension strength 5/5 • Single Leg Squat symmetrical with uninvolved side • Full Pain-free ROM
PHASE IV	20+	<p>EXERCISE PROGRESSION</p> <ul style="list-style-type: none"> • Maintain muscle activation series, trunk, hip and lower extremity strength and flexibility pro-gram • Introduce and progress plyometric program • Begin ladder drills and multidirectional movement • Begin Interval running program • Field/court sports specific drills in controlled environment • Pass sports test • Non-contact drills and scrimmaging – must have passed sports test- refer to specific return to sport program • Return to full activity per physician and passing PT sport test <p>CRITERIA FOR RETURN TO PLAY</p> <ul style="list-style-type: none"> • Continue soft tissue mobilization as needed particularly glutes, adductors, hip flexors, abduc-tors • Gentle joint mobilizations as needed for patients lacking end range FABER ROM • Trigger point dry needling for glutes, TFL, quads, adductors, iliopsoas, iliacus may continue to benefit patients with tightness or mild ROM restrictions 	<ul style="list-style-type: none"> • Return to full activity